



Direction of travel
past pharmacode scanner

First bar

Insert Perrigo
component number here

If you need more advice ask your pharmacist

Active ingredient
Each 5.6 g sachet of granules contains Sodium Citrate Dihydrate 4 g.
Also contains: sucrose, colloidal silicon dioxide, saccharin sodium, cranberry flavour.
PL 12063/0045 UK Only Text prepared 05/24
Manufactured for The Boots Company PLC Nottingham NG2 3AA by the Marketing Authorisation holder Wrafton Laboratories Limited Braunton Devon EX33 2DL



CRANBERRY FLAVOUR
> 48-hour complete course
6 SACHETS
EFFECTIVE RELIEF OF THE SYMPTOMS OF
CYSTITIS INCLUDING BURNING AND STINGING

Boots
Cystitis Relief
4g Granules for Oral Solution
Sodium Citrate

Boots
Cystitis Relief
4g Granules for Oral Solution
Cor Scheme at www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.
Reporting of side effects
If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed on this carton. You can also report side effects directly via the Yellow Card Scheme at www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.
How to store this medicine
Do not store above 25°C. Store in a dry place. Keep out of the sight and reach of children.
Use by the date on the end flap of the carton.

Read all of this carton for full instructions.
What this medicine is for
This medicine contains Sodium Citrate which relieves the symptoms of cystitis in women by making the urine less acidic.
Before you take this medicine
Do not take if you

- are allergic to any of the ingredients
- have diabetes, heart disease, high blood pressure
- have kidney disease or have had it
- are on a low salt (sodium) diet
- have an intolerance to some sugars, unless your doctor tells you to (this medicine contains sucrose)
- are pregnant or breastfeeding

This medicine contains 939 mg sodium (main component of cooking/table salt) in each sachet. This is equivalent to 48 % of the recommended maximum daily dietary intake of sodium for an adult.
Talk to your doctor if you have symptoms:

- which carry on for more than 4 days after you start taking this medicine
- which keep recurring

Talk to your doctor or pharmacist if you are taking:

- Lithium for manic depression
- Methenamine for urinary infections
- Other medicines as these may be affected by taking this medicine

How to take this medicine
Check the sachet is not broken before use. If it is, do not take that sachet. Dissolve the contents of the sachet in a glass of water and drink the solution. This is a 2-day course of treatment.
Adult Women: 1 sachet, 3 times a day for 2 days. **This medicine must not be used by men or children.**
Do not exceed the stated dose. If symptoms persist or worsen after the 2-day course is completed, consult your doctor or pharmacist. Do not repeat the treatment without medical advice.
If you take too many sachets: Talk to a doctor.
Possible side effects
Most people will not have problems, but some may get some. **If you get the following side effect, see your doctor or pharmacist:**

- Passing of water more frequently

Boots
Cystitis Relief
4g Granules for Oral Solution

Direction of travel
past pharmacode scanner

First bar

178

12.7

78

30

78

29.3

228

21.5

92

30

13